

Hand Therapy

PIP Joint Replacement (Prof Imam)

The Surgery

You have had a replacement of the middle joint of your finger (proximal interphalangeal joint/PIP joint) after having had severe changes in the joint. This surgery is predominantly one for pain relief but by adhering to the following guidelines, we hope to get your function as good as possible.

The joint that has been used in your surgery is an 'Osteotec Silastic Hinge Arthroplasty'. This is a constrained joint that means there is a low risk of dislocation and gives you an opportunity to be working on your movement sooner. However, this still needs to be done in a protected way and hand therapy will offer you guidance with this.



Immediately Following Surgery

- You will be in a bulky dressing for a minimum of 2 days following surgery.
- Prof Imam will make a referral via email or via day surgery for hand therapy.
- Hand therapy will be in touch with this appointment the day following your surgery.

2-5 Days Post Op

Attend Hand Therapy for:

- Removal of bulky dressing and wound management
- Fabrication of a resting volar splint with the PIP joint at 0 degrees for constant use
- Advise on management of post-operative swelling
- Advise on isolated joint exercises. You are to remove the splint 5 times a day to complete these exercises

1-2 Weeks Following Surgery

Attend Hand Therapy for:

- Dressing changes if required
- Check volar resting splint

2 Weeks Following Surgery

Attend Hand Therapy for:

- Removal of sutures and scar management if ready
- Check ROM
- Check post-operative swelling and address appropriately

3 Weeks Following Surgery

Attend Hand Therapy for:

- Scar management
- Check ROM
- Check post-operative swelling and address appropriately

4 Weeks Following Surgery

Attend Hand Therapy for:

- If no concerns you can decrease splint use to night and protection only at the discretion of your therapist
- Check ROM
- Check post-operative swelling and address appropriately
- Progress home exercise programme to include thesaspone exercises

6 Weeks Following Surgery

Attend Hand Therapy for:

- If no concerns use of the splint can stop at the discretion of your therapist
- Check ROM
- Check post-operative swelling and address appropriately
- You may be advised how to encourage flexion passively
- You may be provided with a blocking splint for the MCP joint to encourage PIP joint flexion on activity

6-12 Weeks Following Surgery

Hand therapy input will continue if there is a specific clinical need and goals that haven't yet been achieved. In this scenario your home exercises will be progressed appropriately for you depending on how you are presenting.

The above guidelines have been produced for Prof Imam's patients only, with the stated implant. These are just guidelines and all patients are treated individually depending on their presentation on a case by case basis and as guided by the Consultant.

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty. If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Sarah Joyce, Specialist Occupational Therapist

Version: 1

Published: December 2017

Department: Hand Therapy Department

Review: December 2019

We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.



اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

ਜੇ ਤੁਹਾਨੂੰ ਭਰਜਾਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Se precisa de uma tradução por favor contacte: 01932 723553

Ashford Hospital London Road, Ashford, Middlesex, TW15 3AA Tel: **01784 884488**

St. Peter's Hospital Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk